

NOTEWORTHY COLLECTION

WYOMING

LUZULA SUBCAPITATA (Rydb.) H. D. Harr. (JUN-CACEAE).—Albany Co., wet meadow at lake's edge, W side of Lost Lake, Medicine Bow Mountains, 3350 m, 19 Jul 1989, *J. Haines & C. Regan 7602A* (RM); wet meadow by lake and stream, S side of western lobe of Lost Lake, Medicine Bow Mountains, 3335 m, 26 Aug 1991, *J. Haines and M. Hille 10027* (RM).

Previous knowledge. Colorado woodrush, native in the mountains of Colorado, was considered a state endemic (Weber 1990). Kartesz (2003, updated 2012) recorded *Luzula subcapitata* in 15 Colorado counties, with a continuous range from San Juan Co. north through the mountains to Larimer Co., on the Wyoming border. Dorn (2001) did not include *L. subcapitata* in his Wyoming flora.

Significance. A native addition to the flora of Wyoming. The Lost Lake population of *L. subcapitata* is only 40 air km N of Larimer Co., Colorado, constituting a short extension of its known range.

—PETER F. ZIKA, WTU Herbarium, Box 355325, University of Washington, Seattle, WA. 98195-5325. Zikap@comcast.net.

LITERATURE CITED

- DORN, R. D. 2001. Vascular plants of Wyoming, 3rd ed. Mountain West Publishing, Cheyenne, WY.
KARTESZ, J. T. 2003, updated 2012. Floristic synthesis of North America. CD-ROM Version 2.0, Biota of North America Program, Chapel Hill, NC.
WEBER, W. A. 1990. Colorado flora: eastern slope. University Press of Colorado, Niwot, CO.